



Impact of maritime transport emissions on coastal air quality in Europe

Mar Viana
Institute of Environmental Assessment and Water Research
mar.viana@idaea.csic.es

Outline


1. Impact of international shipping on European air quality
 - Tracers and physico-chemical characteristics
 - Impact on ambient PM and on gaseous pollutants
2. Mitigation strategies:
 - Overview
 - Case study: Marmara Sea (Turkey)
3. Conclusions

(1)

Impact of international shipping on European air quality

Rationale


Emissions from the marine transport sector contribute **significantly** to air pollution globally



Increasing emission source:

- Globalization of manufacturing processes
- Increase of global-scale trade
- Relatively, large efforts to reduce other sources (industrial, power generation, etc.)
- More future growth expected

Human health



Climate




Ecosystems



How much of a problem?

- Different approaches used in different countries
- Not yet achieved the goals for protecting human health



Literature Review



Chemical tracers

Well-known **tracers** of combustion based on crude oil:

- V and Ni (>60 publications)
- Others: La, Th, Pb, Zn and SO_4^{2-} (>18 publications)

Where?	PM _x	V/Ni	Reference
Italy	PM ₁₀	3.2±0.8	Mazzei et al. (2008)
	PM _{2.5}	3.2±0.8	Mazzei et al. (2008)
	PM ₁₀	3.2±0.8	Mazzei et al. (2008)
Ship engine		2.3-4.5	Agrawal et al. (2008)
Spain	PM _{2.5}	4-5	Viana et al. (2009)
	PM ₁₀	4-5	Viana et al. (2009)
Spain	PM ₁₀	3	Pandolfi et al. (2011)
	PM _{2.5}	3	Pandolfi et al. (2011)
Europe	PM ₁₀	3-4	Viana et al. (2014)
Europe	PM _{2.5}	3-4	Viana et al. (2014)
Europe	PM ₁₀	2.3-2.5	Alastuey et al. (2016)


Where?	PM _x	Tracer	Value	Reference
Spain	PM ₁₀	V/EC	<2	Viana et al. (2009)
	PM _{2.5}	V/EC	<2	Viana et al. (2009)
Spain	PM ₁₀	La/Ce	0.6-0.8	Pandolfi et al. (2011)
	PM _{2.5}	La/Ce	0.6-0.8	Pandolfi et al. (2011)
Italy	PM ₁₀	soluble V	80%	Becagli et al. (2012)
	PM ₁₀	soluble V	>6 ng/m ³	Becagli et al. (2012)
	PM ₁₀	soluble Ni	80%	Becagli et al. (2012)
	PM ₁₀	non-ss SO_4^{2-} /V	200-400	Becagli et al. (2012)

Tracers are available, BUT:
changing fuels result in changing tracers

Other tracers



Shipping emissions correlate with:

- NO, NO_x, SO₂ and VOCs
- Particle number concentration (N): nucleation episodes (SO₂) (Reche et al., 2011)
- Particle size distribution (80-500 nm; e.g., Masiol et al., 2016)



Primary vs. Secondary particles

In Southern-Europe:



Qinhuangdao (China) (Lang et al., 2017):

Primary = 51% of shipping PM2.5 vs. Secondary = 48%; large seasonal variability

Viana et al. (2014)

Premature deaths/year in Europe:



due to primary particles
(301.000/year)



due to secondary particles
(245.000/year)

More efficient for health to decrease primary PM emissions?

Andersson et al. (2009); Hammingh et al. (2012); Tian et al. (2013); Lang et al. (2017)

References:

- Genoa (Italy): Mazzei et al. (2008)
- Melilla (Spain): Viana et al. (2009)
- Cork (Ireland): Hellebust et al. (2010)
- Algeciras (Spain): Pandolfi et al. (2011)
- Lampedusa (Italy): Becagli et al. (2012)
- Barcelona (Spain): Amato et al. (2009)
- Netherlands, UK, Belgium, Denmark, France, Germany, Sweden, Norway, Luxembourg, Switzerland: Hammingh et al. (2012)

- UK: Hadley et al. (2016)


1-7% PM₁₀
1-20% PM_{2.5}
8-11% PM₁

- PM₁₀
- PM_{2.5}
- PM₁

Comparability?

13-17% PM_{2.5} in China
 Shanghai; Pearl River Delta
Zhao et al. (2013); Tao et al. (2017)

10-70% PM_{2.5} in Western USA, Seattle
Hadley (2017)



References:


- Sweden: Isakson et al. (2001)
- Netherlands, Denmark, Germany, France, Ireland: Hammingh et al. (2012)

● NO₂

● SO₂

7-24% NO₂

Comparability?



Contributions to gases (NO, NO₂, SO₂) > PM, N

Hotelling: contribution to SO₂ < NO & NO₂ due to low-S fuels at berth

Contribution to NO >> NO₂ and provoked local-scale depletion of O₃

Impact of harbour operations

- Knowledge gap! Loading and unloading of vessels, fuelling, etc.
- Studies agree on the **relevance of this impact**:
 - S-Europe: road dust = 26% PM₁₀ in harbours; harbours = **9-12%** urban PM₁₀ (*Pérez et al., 2016*)
 - Los Angeles harbour: vehicular sources + road dust = **54%** of PM_x, vs. shipping < 5% of PM_{2.5} (*Minguillon et al., 2008*)
 - Hotelling, manoeuvring (*Merico et al., 2016*)



(2) Mitigation strategies

Mitigation strategies



**IMO (UN), MARPOL,
SECAs, NECAs**



National regulations




**EU Directive 2005/33/EC on sulphur
emissions from ships**


Technological measures:

- low sulphur fuels
- sulphur scrubbers
- NOx mitigation measures
- liquid natural gas (LNG)
- slow steaming
- soot particle filters...

Mitigation strategies



Velders et al. (2011)



Year 2010 << Years 2000-2006
(50%)

SO₂ emissions reduction:


- >2006: use of low-S fuel due to the SECA regulations in the North Sea
- >2007: MARPOL convention
- >2010: EU directive 2005/33/EC

Mitigation strategies

Directive 2005/33/EC:

- SO₂ concentrations in 3 out of 4 harbours decreased (>2010)
- No decrease was observed in Tunis
- Average decrease SO₂ = 66% (daily)
- No significant changes for NO_x & BC

Schembari et al. (2012)



Sulphur reduction policy in the Baltic Sea SECA (2015):

- for the Baltic Sea only, the latest sulphur regulation is not cost-effective
- Expected annual cost = 465 M€
- Monetized benefit = 105 M€

Annturi et al. (2016)


Case study: ECA in the Marmara Sea




Challenges using BenMAP

Pollutant data: modelled SO_2 , PM_{10} , $\text{PM}_{2.5}$ with CALPUFF


SO_2 before ECA




SO_2 after ECA



PM_{10} before ECA



PM_{10} after ECA




Highly spatially-resolved data


Results: Environmental benefits

Istanbul: Air quality improvement

PM_{2.5} before ECA




PM_{2.5} after ECA




■ Total PMx

■ Ship-sourced PMx

SO₂ before ECA



SO₂ after ECA



■ Total SO₂

■ Ship-sourced SO₂

Results: Health benefits



		East domain (90% confidence intervals)		
Health outcome	Scenario	PM ₁₀	PM _{2.5}	SO ₂
Hospital admissions for respiratory diseases (ICD-10 J00-J99)	Baseline (total burden)	13,000 (4,900 to 20,000)	18,000 (6,800 to 20,000)	1,200 (-830 to 3,200)
	Policy scenario (number avoided)	150 (57 to 230)	330 (125 to 370)	180 (-108 to 460)
	% Change	-1%	-2%	-14%
Hospital admissions for circulatory system diseases (ICD-10 I00-I90)	Baseline (total burden)	4,300 (770 to 7,800)	6,000 (1,900 to 9,700)	1,700 (770 to 2,500)
	Policy scenario (number avoided)	45 (8.1 to 82)	97 (30 to 160)	190 (90 to 290)
	% Change	-1%	-2%	-12%
All-cause mortality (ICD-10 A00-R99)	Baseline (total burden)	120 (50 to 190)	670 (140 to 1,000)	17 (15 to 19)
	Policy scenario (number avoided)	1 (0.4 to 1.6)	13 (2.7 to 19)	2 (1.7 to 2.2)
	% Change	-1%	-2%	-10%

(3) Conclusions

Conclusions & knowledge gaps (1)

- What we know:
 - Nr. studies not large, but increasing
 - Impact on PM_x, NO_x, SO₂, and new particle formation (N)
 - Contribution to PM_x: 1-20% PM_x, with large spatial variability
- What we don't know (so well):
 - Is it more efficient to reduce primary or secondary emissions?
 - Impact of harbour operations & how to mitigate them

Conclusions & knowledge gaps (2)

- Mitigation strategies:
 - Efficient: 50-66% SO₂ reduction, and 2^{ary} PM
 - Cost effective?
- Case study: potential improvements in Istanbul
 - Environmental benefits: 46% to 5% reduction of ship-sourced SO₂
Health benefits: 12-14% decreased hospital admissions due to SO₂;
10% reduced mortality due to SO₂
 - **Overall, beneficial policy from an environmental and health perspective**
- Limitations:
 - Uncertainties in emissions modelling & AQ measurements
 - Need for regionally-specific health impact functions



MINISTERIO
DE ECONOMÍA
Y COMPETITIVIDAD



CSIC
CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS



European Topic Centre
on Air Pollution and
Climate Change Mitigation



vito
vision on technology



centre de recerca
en epidemiologia
ambiental



Thank you for your attention

mar.viana@idaea.csic.es

